



LOKELMA™

(sodium zirconium cyclosilicate)

10 g for oral suspension

YOUR TREATMENT MATTERS

Learn about hyperkalemia and LOKELMA

APPROVED USE

LOKELMA is a prescription medication for the treatment of high levels of potassium in the blood (hyperkalemia) in adults.

IMPORTANT SAFETY INFORMATION FOR LOKELMA

What should I tell my doctor before taking LOKELMA?


Before taking LOKELMA, tell your doctor about all your medical conditions, including

- problems having a bowel movement, including severe constipation, a blockage (obstruction) in your bowel, or dry hard stool that will not pass out of your rectum (impaction)
- problems with your bowels after bowel surgery

Please read Important Safety Information on page 7, and click www.lokelmapi.com for full Prescribing Information for LOKELMA.

▶ What is hyperkalemia?

Hyperkalemia is the name for a condition where potassium levels in your blood are higher than normal.



*LOKELMA does not treat hypokalemia.

WHAT CAUSES HYPERKALEMIA?

For some people, high potassium may be caused by another health condition, such as

- Kidney problems
- High blood pressure
- Diabetes
- Serious heart problems

Hyperkalemia can also be caused by some medications.

Please read **Important Safety Information on page 7**, and click www.lokelmapi.com for full **Prescribing Information for LOKELMA**.

▶ Why LOKELMA?

WHY DID MY DOCTOR CHOOSE LOKELMA FOR ME?

LOKELMA is a prescription medicine for adults with hyperkalemia.

- ✓ **LOKELMA helps bring levels of potassium back down to normal**
- **And with continued treatment, LOKELMA helps keep potassium levels down**

Make a follow-up appointment with your doctor



It's important to have regular appointments with your doctor, so he or she can check on your progress.

IMPORTANT SAFETY INFORMATION FOR LOKELMA

What is the most important thing I should know about LOKELMA?

LOKELMA can cause swelling (edema) caused by fluid retention in your body (such as hands, ankles, feet).

- LOKELMA contains 400 mg of sodium in each 5 g dose. Your doctor may tell you to reduce the sodium in your diet or increase the dose of your diuretic
- Watch for signs of swelling in your body especially if you limit sodium in your diet or are likely to have fluid retention in your body due to heart or kidney problems

 **LOKELMA™**
(sodium zirconium cyclosilicate)
10 g for oral suspension

▶ Taking LOKELMA

Take LOKELMA exactly as prescribed and for as long as your doctor prescribes. This matters—a lot.

RECOMMENDED DOSING

INITIAL TREATMENT

TAKE ONE
10g PACKET 3X/DAY
FOR UP TO 48 HOURS

MAINTENANCE THERAPY

TAKE ONE
10g PACKET 1X/DAY

Your doctor should monitor your potassium levels during treatment and may adjust your dose.

- LOKELMA comes in 2 doses
– 5 g or 10 g white powder in a foil-lined packet

HOW DOES LOKELMA WORK?

LOKELMA is not absorbed in the body—it travels through the GI tract. Along the way, LOKELMA binds to excess potassium in exchange for hydrogen and sodium in the intestines and pulls the potassium out of the body.

Please read Important Safety Information on page 7, and click www.lokelmapi.com for full Prescribing Information for LOKELMA.

▶ Taking LOKELMA

If you have been prescribed other oral medications, you may need to take them 2 hours before or 2 hours after taking LOKELMA as your doctor prescribed.

Tell your doctor or pharmacist about all your medications.

HOW TO TAKE LOKELMA AT HOME

- ✓ Fill a drinking glass with approximately 3 tablespoons of water or more if desired
- ✓ Empty the entire contents of the LOKELMA packet(s) into the glass. Stir well and drink immediately
- ✓ If powder remains in the glass, add water, stir, and drink immediately. Repeat until no powder remains
- ✓ If desired, LOKELMA can be taken with or without food

3 tbsp
of water



LOKELMA is tasteless and odorless

IMPORTANT SAFETY INFORMATION FOR LOKELMA

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LOKELMA may affect how some oral medicines work. Ask your doctor or pharmacist if you need to take LOKELMA 2 hours before or after other oral medicines.

 **LOKELMA™**
(sodium zirconium cyclosilicate)
10 g for oral suspension

▶ Join the My *LOKELMA* Support Program

 **LOKELMA™**
(sodium zirconium cyclosilicate)
for oral suspension

 **My LOKELMA™**
Support Program

The *My LOKELMA Support Program* offers many benefits to those who are starting or already taking LOKELMA:

- Insurance coverage and copay information
- Access support
- Hyperkalemia management and educational information



CALL
1-866-494-8080

- Find out how you could save on future refills
- Join the *My LOKELMA Support Program*

▶ Important Safety Information (continued)

IMPORTANT SAFETY INFORMATION FOR LOKELMA™ (sodium zirconium cyclosilicate) 10 g FOR ORAL SUSPENSION

What is the most important thing I should know about LOKELMA?

LOKELMA can cause swelling (edema) caused by fluid retention in your body (such as hands, ankles, feet).

- LOKELMA contains 400 mg of sodium in each 5 g dose. Your doctor may tell you to reduce the sodium in your diet or increase the dose of your diuretic
- Watch for signs of swelling in your body especially if you limit sodium in your diet or are likely to have fluid retention in your body due to heart or kidney problems

What should I tell my doctor before taking LOKELMA?

Before taking LOKELMA, tell your doctor about all your medical conditions, including

- problems having a bowel movement, including severe constipation, a blockage (obstruction) in your bowel, or dry hard stool that will not pass out of your rectum (impaction)
- problems with your bowels after bowel surgery

What are the most common side effects with LOKELMA?

The most common side effect of LOKELMA is swelling in your body caused by fluid retention. Tell your doctor if you have any side effects that bother you or do not go away.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LOKELMA may affect how some oral medicines work. Ask your doctor or pharmacist if you need to take LOKELMA 2 hours before or after other oral medicines.

Please read Important Safety Information on page 7, and click www.lokelmapi.com for full Prescribing Information for LOKELMA.

 **LOKELMA™**
(sodium zirconium cyclosilicate)
10 g for oral suspension



You may report side effects related to AstraZeneca products by clicking [here](#).

If you cannot afford your medication, AstraZeneca may be able to help. For more information, please visit AstraZeneca-US.com

Please read Important Safety Information on page 7, and click www.lokelmapi.com for full Prescribing Information for LOKELMA.

 **LOKELMA™**
(sodium zirconium cyclosilicate)
10 g for oral suspension